

## RIEMANN INTEGRAL OF A RANDOM FUNCTION AND THE PARABOLIC EQUATION WITH A GENERAL STOCHASTIC MEASURE

UDC 519.21

V. RADCHENKO

ABSTRACT. For stochastic parabolic equation driven by a general stochastic measure, the weak solution is obtained. The integral of a random function in the equation is considered as a limit in probability of Riemann integral sums. Basic properties of such integrals are studied in the paper.

АНОТАЦІЯ. Отримано слабкий розв'язок стохастичного параболічного рівняння, в якому випадковий вплив задано інтегралом за загальною стохастичною мірою. Інтеграл від випадкової функції за дійсною мірою в рівнянні розуміється як границя за ймовірністю інтегральних сум Рімана. Досліджено основні властивості цього інтеграла.

Аннотация. Получено слабое решение стохастического параболического уравнения, в котором случайное влияние задано интегралом по общей стохастической мере. Интеграл от случайной функции по действительной мере в уравнении понимается как предел по вероятности интегральных сум Римана. Изучены основные свойства такого интеграла.

### 1. INTRODUCTION

In this paper we consider the stochastic parabolic equation, which can formally be written as

$$dX(x, t) = AX(x, t) dt + f(x, t) d\mu(t), \quad X(x, 0) = \xi(x), \quad (1.1)$$

where  $(x, t) \in \mathbb{R}^d \times [0, T]$ ,  $A$  is a second-order strongly elliptic differential operator, and  $\mu$  is a general stochastic measure defined on the Borel  $\sigma$ -algebra of  $[0, T]$ . For  $\mu$  we assume  $\sigma$ -additivity in probability only, assumptions for  $A$ ,  $f$  and  $\xi$  are given in Section 6. Equation (1.1) is interpreted in the weak sense (see (6.1) below). We prove existence and uniqueness of solution.

Weak form of (1.1) includes the integral of random function with respect to deterministic measure (Jordan content). We interpret this integral as a limit in probability of Riemann integral sums. This definition of the integral allows to interchange the order of integration with respect to deterministic and stochastic measures (Theorem 4.1), that is important for solving the equation. A large part of the paper is devoted to the study of this Riemann-type stochastic integral.

Parabolic stochastic partial differential equations (SPDEs) driven by the martingale measures had been introduced and discussed initially in [19]. This approach was developed in [1, 3]. Parabolic SPDEs as equations in infinite dimensional space were studied in [4, 11]. In these and many other papers the stochastic noise has some distributional, integrability or martingale properties. In our paper, we consider very general class of

---

2000 *Mathematics Subject Classification.* Primary 60H05, 60H15.

*Key words and phrases.* Stochastic measure, Riemann integral, stochastic parabolic equation, weak solution.

This research was supported by Alexander von Humboldt Foundation, grant no. UKR/1074615. The author wishes to thank Prof. M. Zähle for fruitful discussions, and the hospitality of Jena University is gratefully acknowledged.

possible  $\mu$  on  $[0, T]$ . On the other hand, the stochastic term in (1.1) is independent of  $u$ . A reason is that appropriate definition of integral of random function with respect to  $\mu$  does not exist.

Some motivating examples for studying SPDEs may be found in [4, Introduction], [6, section 13.2]. For  $A = \Delta$ , equation (1.1) describes the evolution in time of the density  $X$  of some quantity such a heat or chemical concentration in a system with random sources. In our model, the random influence can be rather general.

## 2. PRELIMINARIES

Let  $L_0 = L_0(\Omega, \mathcal{F}, \mathbb{P})$  be a set of all real-valued random variables defined on a complete probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  (equivalence classes of). Convergence in  $L_0$  means the convergence in probability and is the convergence in the quasi-norm

$$\|\eta\| = \inf\{\delta: \mathbb{P}\{|\eta| > \delta\} \leq \delta\}.$$

Note that  $\|\eta_1 + \eta_2\| \leq \|\eta_1\| + \|\eta_2\|$ . The following inequality will be used in the sequel

$$\left\| \sum_{k=1}^l c_k \xi_k \right\| \leq 8 \max_{a_k = \pm 1} \left\| \sum_{k=1}^l a_k \xi_k \right\| \leq 16 \max_V \left\| \sum_{k \in V} \xi_k \right\|, \quad |c_k| \leq 1, \quad \xi_k \in L_0, \quad (2.1)$$

where the latter maximum is taken over all possible  $V \subset \{1, \dots, l\}$  (see [16, Theorem 3]).

Let  $\mathcal{S}$  be an arbitrary set and  $\mathcal{B}$  be a  $\sigma$ -algebra of subsets of  $\mathcal{S}$ .

**Definition 2.1.** Any  $\sigma$ -additive mapping  $\mu: \mathcal{B} \rightarrow L_0$  is called a *stochastic measure*.

In other words,  $\mu$  is a vector measure with values in  $L_0$ . We do not assume positivity or integrability for stochastic measures. In [7] such a  $\mu$  is called a general stochastic measure. In the following,  $\mu$  always denotes a stochastic measure.

Examples of stochastic measures are the following. Let  $\mathcal{S} = [0, T] \subset \mathbb{R}_+$ ,  $\mathcal{B}$  be the  $\sigma$ -algebra of Borel subsets of  $[0, T]$ , and  $Y(t)$  be a square integrable martingale. Then  $\mu(\mathbf{A}) = \int_0^T \mathbf{1}_{\mathbf{A}}(t) dY(t)$  is a stochastic measure. If  $W^H(t)$  is a fractional Brownian motion with Hurst index  $H > 1/2$  and  $f: [0, T] \rightarrow \mathbb{R}$  is a bounded measurable function then  $\mu(\mathbf{A}) = \int_0^T f(t) \mathbf{1}_{\mathbf{A}}(t) dW^H(t)$  is also a stochastic measure, as follows from [8, Theorem 1.1]. Some other examples may be found in [7, subsection 7.2]. Theorem 8.3.1 [7] states the conditions under which the increments of a real-valued Lévy process generate a stochastic measure.

For deterministic measurable functions  $g: \mathcal{S} \rightarrow \mathbb{R}$ , an integral of the form  $\int_{\mathcal{S}} g d\mu$  is studied in [12] (see also [7, Chapter 7], [2]). The construction of this integral is standard, uses an approximation by simple functions and is based on results of [15, 17, 18]. In particular, every bounded measurable  $g$  is integrable with respect to any  $\mu$ . An analogue of the Lebesgue dominated convergence theorem holds for this integral (see [7, Proposition 7.1.1] or [12, Corollary 1.2]).

For equations with stochastic measures, weak solutions of some SPDEs were obtained in [13]. Regularity properties of mild solution of the stochastic heat equation were considered in [14].

## 3. RIEMANN INTEGRAL OF A RANDOM FUNCTION

Let  $\mathbf{B} \subset \mathbb{R}^d$  be a Jordan measurable set, and  $\xi: \mathbf{B} \rightarrow L_0$  be a random function. We shall say that  $\xi$  has an integral on  $\mathbf{B}$  if for any sequence of partitions

$$\mathbf{B} = \bigcup_{1 \leq k \leq k_n} \mathbf{B}_{kn}, \quad n \geq 1, \quad \max_k \text{diam } \mathbf{B}_{kn} \rightarrow 0, \quad n \rightarrow \infty, \quad x_{kn} \in \mathbf{B}_{kn},$$

the limit in probability

$$p \lim_{n \rightarrow \infty} \sum_{1 \leq k \leq k_n} \xi(x_{k_n}) m(B_{k_n}) = \int_B \xi(x) dx \tag{3.1}$$

exists. Here  $m$  denotes the Jordan content, sets  $B_{k_n}$ ,  $1 \leq k \leq k_n$ , are assumed to be Jordan measurable and have no common interior points. By mixing of different sequences of partitions, we can prove that the limit is independent of the choice of the sequence. For deterministic  $\xi$ , our definition is equivalent to the definition of the standard Riemann integral in [9].

**Lemma 3.1.** *Let  $\xi$  has an integral on  $B = \prod_{k=1}^d [a_k, b_k] \subset \mathbb{R}^d$ . Then the set of values  $\{\xi(x), x \in B\}$  is bounded in probability.*

*Proof.* Is analogous to the deterministic case. □

For some other  $B \subset \mathbb{R}^d$ , limit (3.1) can exists for unbounded  $\xi$  (for instance, in the case  $m(B) = 0$ ). We use the following

**Definition 3.1.** Random function  $\xi$  is called *integrable on B* if  $\xi$  has an integral on B and set of values  $\{\xi(x), x \in B\}$  is bounded in probability.

Let  $\tilde{B} \subset \mathbb{R}^d$  be an unbounded set for which there exists a sequence of Jordan measurable sets  $B^{(j)}$  such that

$$B^{(j)} \uparrow \tilde{B}, \quad \forall c > 0 \exists j: \tilde{B} \cap \{|x| \leq c\} \subset B^{(j)} \tag{3.2}$$

(we call  $B^{(j)}$  the exhaustive sets). We shall say that  $\xi$  is integrable (in improper sense) on  $\tilde{B}$ , if  $\xi$  is integrable on each  $B^{(j)}$ , and there exists the limit in probability

$$p \lim_{j \rightarrow \infty} \int_{B^{(j)}} \xi(x) dx = \int_{\tilde{B}} \xi(x) dx,$$

that is independent of choice of  $B^{(j)}$ .

All bounded subsets of  $\mathbb{R}^d$  used in the paper are assumed to be Jordan measurable, and all unbounded sets are assumed to be approximable by Jordan measurable sets in the sense of (3.2). Sets in partitions are assumed to be non-overlapping.

Obviously, if  $\xi$  has the Riemann integrable paths then  $\xi$  is integrable in our sense. Theorem 4.1 below gives other examples of integrable random functions.

Further, we establish basic properties of the integral.

**Lemma 3.2.** *Let  $\xi$  be integrable on B. Then  $\xi$  is integrable on each  $A \subset B$ , and for any  $\varepsilon > 0$  there exists  $\delta > 0$  such that for all  $A \subset B$ ,  $A = \bigcup_{1 \leq k \leq k_0} A_k$ ,  $x_k \in A_k$ ,  $\text{diam } A_k < \delta$ , holds*

$$\left\| \sum_{1 \leq k \leq k_0} \xi(x_k) m(A_k) - \int_A \xi(x) dx \right\| < \varepsilon.$$

*Proof.* Suppose the lemma were false. Then

$$\exists \varepsilon_0 > 0 \forall \delta > 0 \exists A = \bigcup_{1 \leq k \leq k_0} A_k = \bigcup_{1 \leq i \leq i_0} A'_i, \quad \text{diam } A_k, \text{diam } A'_i < \delta:$$

$$\left\| \sum_{1 \leq k \leq k_0} \xi(x_k) m(A_k) - \sum_{1 \leq i \leq i_0} \xi(x'_i) m(A'_i) \right\| \geq \varepsilon_0.$$

Take an arbitrary partition

$$B \setminus A = \bigcup_{1 \leq j \leq j_0} C_j, \quad \text{diam } C_j < \delta,$$

and add

$$\sum_{1 \leq j \leq j_0} \xi(x''_j) \mathfrak{m}(C_j), \quad x''_j \in C_j,$$

to each of the considered sums on  $\mathbf{A}$ . Thus we can get two integral sums on  $\mathbf{B}$  with arbitrary small diameters such that the quasi-norm of their difference is greater than or equal to  $\varepsilon_0$ . This contradicts the integrability of  $\xi$  on  $\mathbf{B}$ .  $\square$

**Lemma 3.3.** *Let  $\xi$  be integrable on  $\tilde{\mathbf{B}}$  in the improper sense,  $\tilde{\mathbf{A}} \subset \tilde{\mathbf{B}}$ . Then  $\xi$  is integrable on  $\tilde{\mathbf{A}}$  (if  $\tilde{\mathbf{A}}$  is an unbounded set, the integral is meant in the improper sense).*

*Proof.* Take exhaustive sets  $\mathbf{B}^{(j)} \uparrow \tilde{\mathbf{B}}$ ,  $\mathbf{A}^{(i)} \uparrow \tilde{\mathbf{A}}$ . Then sets

$$(\mathbf{B}^{(j)} \setminus \tilde{\mathbf{A}}) \cup \mathbf{A}^{(i)} \uparrow \tilde{\mathbf{B}}, \quad i, j \rightarrow \infty$$

are exhaustive too, and

$$\int_{\tilde{\mathbf{B}}} \xi(x) dx = \text{p} \lim_{i, j \rightarrow \infty} \left( \int_{\mathbf{B}^{(j)} \setminus \tilde{\mathbf{A}}} \xi(x) dx + \int_{\mathbf{A}^{(i)}} \xi(x) dx \right). \quad (3.3)$$

If  $\text{p} \lim_{i \rightarrow \infty} \int_{\mathbf{A}^{(i)}} \xi(x) dx$  does not exist then we can choose  $i, j \rightarrow \infty$  such that the limit in (3.3) does not exist.  $\square$

**Lemma 3.4.** *Let  $\xi$  be integrable on  $\mathbf{B}$ . Then the set of values  $\{\int_{\mathbf{A}} \xi(s) ds, \mathbf{A} \subset \mathbf{B}\}$  is bounded in probability.*

*Proof.* Suppose the lemma were false. Then

$$\exists \varepsilon_0 > 0, \mathbf{A}_n \subset \mathbf{B}, n \geq 1: \left\| \frac{1}{n} \int_{\mathbf{A}_n} \xi(s) ds \right\| \geq \varepsilon_0.$$

By Lemma 3.2, we can choose a partition  $\mathbf{B} = \bigcup_{1 \leq k \leq k_0} \mathbf{B}_k$  fine enough, such that all integral sums for partitions  $\mathbf{A}_n = \bigcup_{1 \leq k \leq k_0} (\mathbf{A}_n \cap \mathbf{B}_k)$  will be close enough to the integrals on  $\mathbf{A}_n$ . Thus, for all  $n$ ,  $x_{kn} \in \mathbf{A}_n \cap \mathbf{B}_k$ , we get

$$\left\| \frac{1}{n} \sum_{1 \leq k \leq k_0} \xi(x_{kn}) \mathfrak{m}(\mathbf{A}_n \cap \mathbf{B}_k) \right\| \geq \frac{\varepsilon_0}{2}.$$

Since the number of summands is fixed for all  $n$ , we arrive at a contradiction with boundedness of  $\xi$ .  $\square$

**Lemma 3.5.** *Let  $\xi$  be integrable on  $\mathbf{B}$ ,  $f: \mathbf{B} \rightarrow \mathbb{R}$  be a deterministic uniformly continuous on  $\mathbf{B}$  function. Then  $f\xi$  is integrable on  $\mathbf{B}$ .*

*Proof.* Consider the difference of two integral sums of  $f\xi$

$$\begin{aligned} & \left\| \sum_{1 \leq k \leq k_m} f(x_{km}) \xi(x_{km}) \mathfrak{m}(\mathbf{B}_{km}) - \sum_{1 \leq i \leq i_n} f(x_{in}) \xi(x_{in}) \mathfrak{m}(\mathbf{B}_{in}) \right\| \\ &= \left\| \sum_{1 \leq k \leq k_m, 1 \leq i \leq i_n} [f(x_{km}) \xi(x_{km}) - f(x_{in}) \xi(x_{in})] \mathfrak{m}(\mathbf{B}_{km} \cap \mathbf{B}_{in}) \right\| \\ &\leq \left\| \sum_{1 \leq k \leq k_m, 1 \leq i \leq i_n} [\xi(x_{km}) - \xi(x_{in})] f(x_{in}) \mathfrak{m}(\mathbf{B}_{km} \cap \mathbf{B}_{in}) \right\| \\ &\quad + \left\| \sum_{1 \leq k \leq k_m, 1 \leq i \leq i_n} [f(x_{km}) - f(x_{in})] \xi(x_{km}) \mathfrak{m}(\mathbf{B}_{km} \cap \mathbf{B}_{in}) \right\| \\ &= S_1 + S_2. \end{aligned}$$

From (2.1) for  $|f(x)| \leq C$  we get

$$S_1 \leq 16 \max_V \left\| C \sum_{(k,i) \in V} [\xi(x_{km}) - \xi(x_{in})] m(\mathbf{B}_{km} \cap \mathbf{B}_{in}) \right\|, \quad (3.4)$$

where the maximum is taken over all possible sets of pairs  $(k, i)$ .

For example, consider

$$\begin{aligned} & \sum_{(k,i) \in V} \xi(x_{km}) m(\mathbf{B}_{km} \cap \mathbf{B}_{in}) \\ &= \sum_{1 \leq k \leq k_m} \xi(x_{km}) \left[ \sum_{i: (k,i) \in V} m(\mathbf{B}_{km} \cap \mathbf{B}_{in}) + m(\mathbf{B}_{km} \cap \mathbf{B}_{i'n}) \mathbf{1}_{x_{km} \notin (\cup_{i: (k,i) \in V} \mathbf{B}_{in})} \right] \\ & \quad - \sum_{1 \leq k \leq k_m} \xi(x_{km}) m(\mathbf{B}_{km} \cap \mathbf{B}_{i'n}) \mathbf{1}_{x_{km} \notin (\cup_{i: (k,i) \in V} \mathbf{B}_{in})} \\ &= I_1 - I_2. \end{aligned}$$

Here  $\mathbf{B}_{i'n}$  is one of the sets  $\mathbf{B}_{in}$ ,  $1 \leq i \leq i_n$ , that contains  $x_{km}$ . (If  $x_{km}$  lies on the border of  $\mathbf{B}_{i'n}$ , we take it only once.)  $I_1$  and  $I_2$  are integral sums and, by Lemma 3.2, they approximate the integrals of  $\xi$  on respective sets. Therefore, for diameter small enough,  $I_1 - I_2$  will be close to the integral on  $\bigcup_{(k,i) \in V} (\mathbf{B}_{km} \cap \mathbf{B}_{in})$ . Similarly,

$$\sum_{(k,i) \in V} \xi(x_{in}) m(\mathbf{B}_{km} \cap \mathbf{B}_{in})$$

approximate the integral on the same set, and we make the right hand side of (3.4) arbitrary small by choosing the diameter.

Further, for any  $\alpha > 0$ , for diameter small enough and  $\mathbf{B}_{km} \cap \mathbf{B}_{in} = \emptyset$ , we have  $|f(x_{km}) - f(x_{in})| < \alpha$  in  $S_2$ . Inequality (2.1) implies

$$S_2 \leq 16 \max_V \left\| \alpha \sum_{(k,i) \in V} \xi(x_{km}) m(\mathbf{B}_{km} \cap \mathbf{B}_{in}) \right\|.$$

As before, we can make the sum arbitrary close to the integral on  $\bigcup_{(k,i) \in V} (\mathbf{B}_{km} \cap \mathbf{B}_{in})$ . From Lemma 3.4 it follows that  $S_2 \rightarrow 0$  as  $\alpha \rightarrow 0$ .  $\square$

**Lemma 3.6.** *Let  $\xi$  be integrable on  $\mathbf{B}$ ,  $f: \mathbf{B} \rightarrow \mathbb{R}$  be a deterministic uniformly continuous on  $\mathbf{B}$  function,  $|f(x)| \leq C$ . Then*

$$\left\| \int_{\mathbf{B}} f(x) \xi(x) dx \right\| \leq 16 \sup_{A \subset \mathbf{B}} \left\| C \int_A \xi(x) dx \right\|.$$

*Proof.* The inequality for respective integral sums follows from (2.1). Further, we pass to the limit and apply Lemmas 3.2 and 3.5.  $\square$

**Lemma 3.7.** *Let  $\xi$  be integrable on  $\mathbf{B}$ ,  $f_n: \mathbf{B} \rightarrow \mathbb{R}$ ,  $n \geq 1$ , be a deterministic uniformly continuous on  $\mathbf{B}$  functions,  $\sup_{x \in \mathbf{B}} |f_n(x)| \rightarrow 0$ ,  $n \rightarrow \infty$ . Then*

$$\int_{\mathbf{B}} f_n(x) \xi(x) dx \xrightarrow{P} 0, \quad n \rightarrow \infty.$$

*Proof.* The statement follows from Lemmas 3.4 and 3.6.  $\square$

**Lemma 3.8.** *Let  $\xi$  be integrable on an unbounded set  $\tilde{\mathbf{B}}$  in improper sense,  $f: \tilde{\mathbf{B}} \rightarrow \mathbb{R}$  be a deterministic bounded uniformly continuous on  $\tilde{\mathbf{B}}$  function. Then  $f\xi$  is integrable on  $\tilde{\mathbf{B}}$  in improper sense.*

*Proof.* For  $B^{(j)} \uparrow \tilde{B}$  and  $|f(x)| \leq C$  Lemma 3.6 implies

$$\left\| \int_{B^{(j)} \setminus B^{(i)}} f(x)\xi(x) dx \right\| \leq 16 \sup_{A \subset (B^{(j)} \setminus B^{(i)})} \left\| C \int_A \xi(x) dx \right\|. \tag{3.5}$$

If the left hand side of (3.5) does not tend to 0 as  $i, j \rightarrow \infty$ , then we can construct a sequence of bounded sets  $C^{(j)} \uparrow \tilde{B}$  such that the sequence  $\int_{C_j} \xi(x) dx, j \geq 1$ , is, non-fundamental.  $\square$

**Lemma 3.9.** *Let  $\xi$  be integrable on unbounded set  $\tilde{B}$  in improper sense,  $f_n: \tilde{B} \rightarrow \mathbb{R}$  be a deterministic bounded uniformly continuous on  $\tilde{B}$  functions,  $\sup_{n \geq 1, x \in \tilde{B}} |f_n(x)| = C < \infty, \sup_{x \in B} |f_n(x)| \rightarrow 0, n \rightarrow \infty$ , for all bounded  $B \subset \tilde{B}$ . Then*

$$\int_{\tilde{B}} f_n(x)\xi(x) dx \xrightarrow{P} 0, \quad n \rightarrow \infty.$$

*Proof.* Suppose the lemma is false. Applying Lemma 3.7, one can find  $\varepsilon_0 > 0$ , subsequence  $f_{n_j}, j \geq 1$ , and bounded disjoint sets  $B_j \subset (\tilde{B} \cap \{|x| \geq j\})$  such that

$$\left\| \int_{B_j} f_{n_j}(x)\xi(x) dx \right\| > \varepsilon_0.$$

From Lemma 3.6 it follows that there exist bounded disjoint sets  $A_j \subset (\tilde{B} \cap \{|x| \geq j\})$  such that  $\|C \int_{A_j} \xi(x) dx\| > (\varepsilon_0/16)$ . This contradicts the integrability of  $\xi$  on  $\tilde{B}$ .  $\square$

Note that the stochastic continuity of  $\xi$  does not imply the integrability.

**Example 3.1.** Consider  $B = [0, 1], \xi_k(\omega) = 5^k \mathbf{1}_{F_k}, k \geq 1$ , where  $P(F_k) = 1/k, F_k$  are independent. Set

$$\begin{aligned} \xi(0) &= 0, \quad \xi(x) = \xi_k, 2^{-2k-1} \leq x \leq 2^{-2k}, \\ \xi(x) &= 2^{2k+2}((2^{-2k-1} - x)\xi_{k+1} + (x - 2^{-2k-2})\xi_k), 2^{-2k-2} \leq x \leq 2^{-2k-1}. \end{aligned}$$

Taking all possible finite unions  $A = \bigcup_k [2^{-2k-1}, 2^{-2k}]$ , we see that the values  $\int_A \xi(x) dx$  are not bounded in probability. By Lemma 3.4,  $\xi$  is not integrable on  $[0, 1]$ .

#### 4. INTERCHANGE OF THE ORDER OF INTEGRATION

**Theorem 4.1.** *Let  $\mu$  be a stochastic measure on  $(S, \mathcal{B}), B \subset \mathbb{R}^d$  be a bounded set. Assume that  $h(x, s): B \times S \rightarrow \mathbb{R}$  is a measurable deterministic function which is Riemann integrable on  $B$  for each fixed  $s$ , and  $|h(x, s)| \leq g(s)$ , where  $g: S \rightarrow \mathbb{R}$  is integrable on  $S$  with respect to  $d\mu(s)$ . Then the random function  $\xi(x) = \int_S h(x, s) d\mu(s)$  is integrable on  $B$ , and*

$$\int_B dx \int_S h(x, s) d\mu(s) = \int_S d\mu(s) \int_B h(x, s) dx. \tag{4.1}$$

*Proof.* From the inequality  $|h(x, s)| \leq g(s)$  and (2.1) it follows that values of  $\xi$  are bounded in probability (see Lemma 1.1 and Theorem 1.3 [12]). Integral sums of  $\int_B \xi(x) dx$  have the form

$$\begin{aligned} \sum_{1 \leq k \leq k_n} m(B_{kn}) \int_S h(x_{kn}, s) d\mu(s) &= \int_S g_n(s) d\mu(s), \\ g_n(s) &= \sum_{1 \leq k \leq k_n} h(x_{kn}, s) m(B_{kn}) \rightarrow \int_B h(x, s) dx. \end{aligned}$$

Boundedness condition of  $h$  and the analogue of the Lebesgue theorem [7, Proposition 7.1.1] for the integral with respect to  $d\mu(s)$  imply the statement.  $\square$

**Corollary 4.1.** *Let  $\mu$  be a stochastic measure on  $(S, \mathcal{B})$ ,  $\tilde{B} \subset \mathbb{R}^d$  be an unbounded set. Assume that  $h(x, s): \tilde{B} \times S \rightarrow \mathbb{R}$  is a measurable deterministic function which is Riemann integrable on  $\tilde{B}$  in improper sense for each fixed  $s$ , and  $|h(x, s)| \leq g(s)$ ,  $\int_{\tilde{B}} |h(x, s)| dx = g_1(s)$ , where  $g, g_1: S \rightarrow \mathbb{R}$  are integrable on  $S$  with respect to  $d\mu(s)$ . Then the random function  $\xi(x) = \int_S h(x, s) d\mu(s)$  is integrable on  $\tilde{B}$  in improper sense, and*

$$\int_{\tilde{B}} dx \int_S h(x, s) d\mu(s) = \int_S d\mu(s) \int_{\tilde{B}} h(x, s) dx. \quad (4.2)$$

*Proof.* For bounded sets  $B^{(j)} \uparrow \tilde{B}$ , Theorem 4.1 implies

$$\int_{B^{(j)}} dx \int_S h(x, s) d\mu(s) = \int_S d\mu(s) \int_{B^{(j)}} h(x, s) dx.$$

Further, we use the analogue of the Lebesgue theorem and integrability of  $g_1$ .  $\square$

**Theorem 4.2.** *Let  $B \subset \mathbb{R}^d$ ,  $S \subset \mathbb{R}^m$  be a bounded sets, random function  $\xi(x, s): B \times S \rightarrow L_0$  be integrable on  $B \times S$  with respect to  $dx \times ds$  and be integrable on  $S$  with respect to  $ds$  for each fixed  $x$ . Then*

$$\int_{B \times S} \xi(x, s) dx \times ds = \int_B dx \int_S \xi(x, s) ds. \quad (4.3)$$

*Proof.* Integral sums of integral with respect to  $dx$  in (4.3) has the form

$$\sum_{1 \leq k \leq k_0} m(B_k) \int_S \xi(x_k, s) ds. \quad (4.4)$$

Each integral in (4.4) may be approximated by sums of the form  $\sum_{1 \leq i \leq i_0} m(S_i) \xi(x_k, s_i)$ . Thus, the sums

$$\sum_{1 \leq k \leq k_0} \sum_{1 \leq i \leq i_0} m(B_k) m(S_i) \xi(x_k, s_i).$$

will approximate the right hand side of (4.4). But they are the integral sums for the integral with respect to  $dx \times ds$  in (4.3), and will be close to the left hand side of (4.3) for sufficiently small diameters of  $B_k \times S_i$ .  $\square$

**Corollary 4.2.** *Let  $S \subset \mathbb{R}^m$  be a bounded set,  $\tilde{B} \subset \mathbb{R}^d$  be an unbounded set. Assume that the random function  $\xi(x, s): \tilde{B} \times S \rightarrow L_0$  is integrable on  $\tilde{B} \times S$  with respect to  $dx \times ds$  in improper sense, is integrable on  $\tilde{B}$  with respect to  $dx$  in improper sense for each fixed  $s$ , and is integrable on  $S$  with respect to  $ds$  for each fixed  $x$ . Then*

$$\int_{\tilde{B} \times S} \xi(x, s) dx \times ds = \int_S ds \int_{\tilde{B}} \xi(x, s) dx = \int_{\tilde{B}} dx \int_S \xi(x, s) ds. \quad (4.5)$$

*Proof.* Consider exhaustive sets  $B^{(j)} \uparrow \tilde{B}$ . For the first of the repeated integrals (4.5), the integral sums has the form

$$\sum_{1 \leq i \leq i_0} m(S_i) \int_{\tilde{B}} \xi(x, s_i) dx \quad (4.6)$$

The integrals in (4.6) can be approximated by  $\int_{B^{(j)}} \xi(x, s_i) dx$ , and the last integral is the limit of sums

$$\sum_{1 \leq k \leq k_0} m(B_k^{(j)}) \xi(x_k^{(j)}, s_i).$$

If integral sums (4.6) does not converge, then we can construct a non-convergent sequence of sums

$$\sum_{1 \leq i \leq i_0} \sum_{1 \leq k \leq k_0} m(S_i) m(B_k^{(j)}) \xi(x_k^{(j)}, s_i),$$

and this contradicts the integrability of  $\xi$  on  $\mathbb{S} \times \tilde{\mathbb{B}}$ .

Further, by Theorem 4.2, for each  $j$  we have

$$\int_{\mathbb{B}^{(j)} \times \mathbb{S}} \xi(x, s) dx \times ds = \int_{\mathbb{B}^{(j)}} dx \int_{\mathbb{S}} \xi(x, s) ds.$$

The left hand side has the limit in probability as  $j \rightarrow \infty$ . Hence, the right hand side has the limit, and the second equality of (4.5) holds.  $\square$

## 5. INTEGRATION BY PARTS

To solve the parabolic stochastic equation, we need the following two lemmas.

**Lemma 5.1.** *Let a random function  $\xi(u): [0, s] \rightarrow L_0$  be integrable on  $[0, s]$ . Then  $\eta(u) = \int_0^u \xi(v) dv$  is integrable on  $[0, s]$ , and*

$$\int_0^s du \int_0^u \xi(v) dv = \int_0^s (s-v)\xi(v) dv.$$

*Proof.* By Lemma 3.5, the function  $(s-v)\xi(v)$  is integrable, by Lemma 3.2  $\eta(u)$  is well defined. The integral sum of  $\int_0^s \eta(u) du$  has the form

$$\sum_{1 \leq k \leq k_0} m(\mathbb{B}_k) \int_0^{u_k} \xi(v) dv, \quad u_k \in \mathbb{B}_k. \quad (5.1)$$

We can take a new partition  $[0, s] = \bigcup_{1 \leq i \leq i_0} \mathbb{C}_i$  such that each integral  $\int_0^{u_k} \xi(v) dv$  be close enough to integral sum with this partition (Lemma 3.2). Thus we can approximate (5.1) arbitrary closely by the sum

$$\sum_{1 \leq k \leq k_0} m(\mathbb{B}_k) \sum_{1 \leq i \leq i_0} m(\mathbb{C}_i \cap [0, u_k]) \xi(v_i), \quad v_i \in \mathbb{C}_i. \quad (5.2)$$

For  $\int_0^s (s-v)\xi(v) dv$ , take the integral sum

$$\sum_{1 \leq i \leq i_0} m(\mathbb{C}_i) (s-v_i) \xi(v_i). \quad (5.3)$$

The difference of (5.3) and (5.2) is equal to

$$\sum_{1 \leq i \leq i_0} \xi(v_i) [m(\mathbb{C}_i)(s-v_i) - m(\mathbb{C}_i) \sum_{k: \mathbb{C}_i < \mathbb{B}_k} m(\mathbb{B}_k) - \sum_{k: \mathbb{C}_i \cap \mathbb{B}_k \neq \emptyset} m(\mathbb{B}_k) m(\mathbb{C}_i \cap [0, u_k])]. \quad (5.4)$$

Notation  $\mathbb{C}_i < \mathbb{B}_k$  means that  $v < u$  for all  $v \in \mathbb{C}_i, u \in \mathbb{B}_k$ . We have

$$0 \leq (s-v_i) - \sum_{k: \mathbb{C}_i < \mathbb{B}_k} m(\mathbb{B}_k) \leq \max_i \text{diam } \mathbb{C}_i + \max_k m(\mathbb{B}_k).$$

The last sum of (5.4) is not greater than

$$m(\mathbb{C}_i) \sum_{k: \mathbb{C}_i \cap \mathbb{B}_k \neq \emptyset} m(\mathbb{B}_k) \leq m(\mathbb{C}_i) \left( \max_i \text{diam } \mathbb{C}_i + 2 \max_k m(\mathbb{B}_k) \right).$$

Therefore, value (5.4) may be written in the form  $\sum_{1 \leq i \leq i_0} \xi(v_i) m(\mathbb{C}_i) \alpha_i$ , where  $\alpha_i \rightarrow 0$  as  $\text{diam } \mathbb{C}_i, \text{diam } \mathbb{B}_k \rightarrow 0$ . From (2.1) we obtain

$$\left\| \sum_{1 \leq i \leq i_0} \xi(v_i) m(\mathbb{C}_i) \alpha_i \right\| \leq 16 \max_V \left\| \max_i |\alpha_i| \sum_{i \in V} \xi(v_i) m(\mathbb{C}_i) \right\|. \quad (5.5)$$

The sums  $\sum_{i \in V} \xi(v_i) m(\mathbb{C}_i)$  are close to respective integrals for  $\text{diam } \mathbb{C}_i$  small enough (Lemma 3.2) and values of integrals are bounded in probability (Lemma 3.4). Therefore, the left hand side of (5.5) tends to zero as  $\max_i |\alpha_i| \rightarrow 0$ .  $\square$



**Lemma 5.2.** *Let a random function  $\xi(u): [0, s] \rightarrow L_0$  be integrable on  $[0, s]$ ,  $f \in \mathbb{C}^{(1)}([0, s])$  be a deterministic function. Then*

$$f(s) \int_0^s \xi(u) du = \int_0^s f(u) \xi(u) du + \int_0^s f'(u) du \int_0^u \xi(v) dv. \quad (5.6)$$

*Proof.* From Lemmas 3.5 and 5.1 it follows that the random functions  $\zeta_1(u) = f(u)\xi(u)$ ,  $\zeta_2(u) = f'(u) \int_0^u \xi(v) dv$  are integrable on  $[0, s]$ . First, let us show that for  $0 = u_0 < u_1 < \dots < u_{k_0} = s$ ,  $\alpha = \max_k |u_k - u_{k-1}|$ , we have

$$\sum_{1 \leq k \leq k_0} (f(u_k) - f(u_{k-1})) \int_0^{u_k} \xi(v) dv \xrightarrow{\mathbb{P}} \int_0^s f'(u) du \int_0^u \xi(v) dv, \quad \alpha \rightarrow 0. \quad (5.7)$$

Applying the Lagrange formula and integrability of  $\zeta_2$ , for some  $\tilde{u}_k \in (u_{k-1}, u_k)$  we obtain

$$\begin{aligned} \sum_{1 \leq k \leq k_0} (f(u_k) - f(u_{k-1})) \int_0^{u_k} \xi(v) dv &= \sum_{1 \leq k \leq k_0} f'(\tilde{u}_k)(u_k - u_{k-1}) \int_0^{u_k} \xi(v) dv \\ &= \sum_{1 \leq k \leq k_0} f'(\tilde{u}_k)(u_k - u_{k-1}) \int_0^{\tilde{u}_k} \xi(v) dv + \sum_{1 \leq k \leq k_0} f'(\tilde{u}_k)(u_k - u_{k-1}) \int_{\tilde{u}_k}^{u_k} \xi(v) dv, \\ \sum_{1 \leq k \leq k_0} f'(\tilde{u}_k)(u_k - u_{k-1}) \int_0^{\tilde{u}_k} \xi(v) dv &\xrightarrow{\mathbb{P}} \int_0^s f'(u) du \int_0^u \xi(v) dv, \quad \alpha \rightarrow 0. \end{aligned}$$

For  $C_1 = \max_u |f'(u)|$ , from (2.1) we have

$$\begin{aligned} &\left\| \sum_{1 \leq k \leq k_0} f'(\tilde{u}_k)(u_k - u_{k-1}) \int_{\tilde{u}_k}^{u_k} \xi(v) dv \right\| \\ &\leq 16 \max_V \left\| C_1 \alpha \sum_{k \in V} \int_{\tilde{u}_k}^{u_k} \xi(v) dv \right\| \leq 16 \sup_A \left\| C_1 \alpha \int_A \xi(v) dv \right\|. \end{aligned}$$

From Lemma 3.4 it follows that the last value tends to 0 as  $\alpha \rightarrow 0$ . Therefore, (5.7) is proved.

Integrability of  $\zeta_1$  implies

$$\begin{aligned} \sum_{1 \leq k \leq k_0} f(u_{k-1}) \int_{u_{k-1}}^{u_k} \xi(v) dv &= \sum_{1 \leq k \leq k_0} f(u_{k-1}) \xi(u_{k-1})(u_k - u_{k-1}) \\ &\quad + \sum_{1 \leq k \leq k_0} f(u_{k-1}) \int_{u_{k-1}}^{u_k} (\xi(v) - \xi(u_{k-1})) dv, \\ \sum_{1 \leq k \leq k_0} f(u_{k-1}) \xi(u_{k-1})(u_k - u_{k-1}) &\xrightarrow{\mathbb{P}} \int_0^s f(u) \xi(u) du, \quad \alpha \rightarrow 0. \end{aligned}$$

For  $C_0 = \max_u |f(u)|$ , from (2.1) we get

$$\begin{aligned} &\left\| \sum_{1 \leq k \leq k_0} f(u_{k-1}) \int_{u_{k-1}}^{u_k} (\xi(v) - \xi(u_{k-1})) dv \right\| \leq 16 \max_V \left\| C_0 \sum_{k \in V} \int_{u_{k-1}}^{u_k} (\xi(v) - \xi(u_{k-1})) dv \right\| \\ &= 16 \max_V \left\| C_0 \left( \int_{\cup_{k \in V} [u_{k-1}, u_k]} \xi(v) dv - \sum_{k \in V} \xi(u_{k-1})(u_k - u_{k-1}) \right) \right\|. \end{aligned}$$

By Lemma 3.2, the last value tends to 0 as  $\alpha \rightarrow 0$ .

Further, we take the obvious equality

$$f(s) \int_0^s \xi(v) dv = \sum_{1 \leq k \leq k_0} (f(u_k) - f(u_{k-1})) \int_0^{u_k} \xi(v) dv + \sum_{1 \leq k \leq k_0} f(u_{k-1}) \int_{u_{k-1}}^{u_k} \xi(v) dv$$

and pass to the limit as  $\alpha \rightarrow 0$ .  $\square$

## 6. PARABOLIC EQUATION WITH A GENERAL STOCHASTIC MEASURE

Consider the differential operator

$$Ag(x) = \sum_{1 \leq i, j \leq d} a_{ij}(x) \frac{\partial^2 g(x)}{\partial x_i \partial x_j} + \sum_{1 \leq i \leq d} b_i(x) \frac{\partial g(x)}{\partial x_i} + c(x)g(x),$$

where  $g : \mathbb{R}^d \rightarrow \mathbb{R}$  and  $a_{ij} = a_{ji}$ . Suppose that  $A$  is strongly elliptic in  $\mathbb{R}^d$  (see (4.5) [5]).

**Assumption 1.** All functions  $a_{ij}$ ,  $b_i$ ,  $c$ ,  $\frac{\partial a_{ij}}{\partial x_i}$ ,  $\frac{\partial^2 a_{ij}}{\partial x_i \partial x_j}$ ,  $\frac{\partial b_i}{\partial x_i}$  are bounded and Hölder continuous in  $\mathbb{R}^d$ .

From now on let  $\mu$  be a stochastic measure on Borel subsets of  $[0, T]$ .

We will study the equation

$$dX(x, t) = AX(x, t) dt + f(x, t) d\mu(t), \quad X(x, 0) = \xi(x), \quad (1.1)$$

where  $X : \mathbb{R}^d \times [0, T] \rightarrow L_0$  is an unknown random function.

We consider (1.1) in the weak sense, i.e.

$$\begin{aligned} \int_{\mathbb{R}^d} X(x, t) \varphi(x) dx &= \int_{\mathbb{R}^d} \xi(x) \varphi(x) dx + \int_{\mathbb{R}^d} A^* \varphi(x) dx \int_0^t X(x, s) ds \\ &+ \int_{[0, t]} d\mu(s) \int_{\mathbb{R}^d} f(x, s) \varphi(x) dx \end{aligned} \quad (6.1)$$

for all test functions  $\varphi \in \mathcal{S}(\mathbb{R}^d)$  (rapidly decreasing Schwartz functions from  $\mathbb{C}^\infty(\mathbb{R}^d)$ ). For each fixed  $t \in [0, T]$  equality (6.1) holds a.s. Integrals of random functions with respect to  $dx$  and  $ds$  are considered in Riemann sense (see section 3), and  $A^*$  denotes the adjoint operator of  $A$ .

**Assumption 2.**  $\xi : \mathbb{R}^d \rightarrow L_0$  is such that  $\xi(\cdot, \omega)$  is continuous and bounded in  $\mathbb{R}^d$  for each fixed  $\omega \in \Omega$ .

**Assumption 3.**  $f : \mathbb{R}^d \times [0, T] \rightarrow \mathbb{R}$  is Borel measurable,  $\sup_t |x|^{-k} |f(x, t)| \rightarrow 0$ ,  $|x| \rightarrow \infty$ , for some  $k > 0$ ,  $f(x, \cdot)$  is continuous and bounded in  $\mathbb{R}^d$  for each fixed  $t \in [0, T]$ .

By Theorem 1 §4 [5], under Assumption 1, the equation  $\partial g / \partial t = Ag$  has a fundamental solution  $p(x, y, t - s)$  (recall that coefficients of  $A$  do not depend on  $t$ ). The following estimate is well known:

$$|p(x, y, t)| \leq C_1 t^{-d/2} \exp\{-C_2 |x - y|^2 / t\}$$

(see, for example, (4.16) [5]). Consider the semigroup

$$S(t)g(x) = \int_{\mathbb{R}^d} p(x, y, t)g(y) dy, \quad t > 0, \quad S(0)g(x) = g(x).$$

Theorem 2 §4 [5] implies that for any continuous bounded  $g$

$$S(t)g(x) = g(x) + A \int_0^t [S(s)g(x)] ds. \quad (6.2)$$

**Theorem 6.1.** *Suppose Assumptions 1–3 hold. Then the random function*

$$X(x, t) = S(t)\xi(x) + \int_{[0,t]} [S(t-s)f(x, s)] d\mu(s) \quad (6.3)$$

is the solution of (6.1).

In addition, suppose the operator  $A$  is self-adjoint,  $X(x, t)$  satisfies (6.1), is integrable on  $\mathbb{R}^d \times [0, T]$  with respect to  $dx \times dt$ , is integrable on  $\mathbb{R}^d$  with respect to  $dx$  for each fixed  $t$ , and is integrable on  $[0, T]$  with respect to  $dt$  for each fixed  $x$ . Then  $X(x, t)$  is given by (6.3).

*Proof.* From (6.2) it follows that for  $X_1(x, t) = S(t)\xi(x)$  and  $f = 0$  equality (6.1) holds. For  $X_2(x, t) = \int_{[0,t]} [S(t-s)f(x, s)] d\mu(s)$  we have

$$\begin{aligned} & \int_{\mathbb{R}^d} A^* \varphi(x) dx \int_0^t X_2(s) ds + \int_{[0,t]} d\mu(s) \int_{\mathbb{R}^d} f(x, s) \varphi(x) dx \\ &= \int_{\mathbb{R}^d} A^* \varphi(x) dx \int_0^t ds \int_{[0,s]} [S(s-u)f(x, u)] d\mu(u) \\ & \quad + \int_{[0,t]} d\mu(s) \int_{\mathbb{R}^d} f(x, s) \varphi(x) dx \\ &\stackrel{(4.1)}{=} \int_{\mathbb{R}^d} A^* \varphi(x) dx \int_{[0,t]} d\mu(u) \int_u^t [S(s-u)f(x, u)] ds \\ & \quad + \int_{[0,t]} d\mu(s) \int_{\mathbb{R}^d} f(x, s) \varphi(x) dx \\ &\stackrel{(4.2)}{=} \int_{[0,t]} d\mu(u) \int_{\mathbb{R}^d} A^* \varphi(x) dx \int_u^t [S(s-u)f(x, u)] ds \\ & \quad + \int_{[0,t]} d\mu(s) \int_{\mathbb{R}^d} f(x, s) \varphi(x) dx \\ &= \int_{[0,t]} d\mu(u) \int_{\mathbb{R}^d} \varphi(x) dx A \int_u^t [S(s-u)f(x, u)] ds + \int_{[0,t]} d\mu(s) \int_{\mathbb{R}^d} f(x, s) \varphi(x) dx \\ &\stackrel{(6.2)}{=} \int_{[0,t]} d\mu(u) \int_{\mathbb{R}^d} \varphi(x) dx ([S(t-u)f(x, u)] - f(x, u)) \\ & \quad + \int_{[0,t]} d\mu(s) \int_{\mathbb{R}^d} f(x, s) \varphi(x) dx \\ &= \int_{[0,t]} d\mu(u) \int_{\mathbb{R}^d} \varphi(x) [S(t-u)f(x, u)] dx \\ &\stackrel{(4.2)}{=} \int_{\mathbb{R}^d} \varphi(x) dx \int_{[0,t]} [S(t-s)f(x, s)] d\mu(s) \\ &= \int_{\mathbb{R}^d} X_2(x, t) \varphi(x) dx. \end{aligned}$$

Therefore (6.1) holds for  $X = X_1 + X_2$ .

Finally, we will prove the uniqueness of the solution. Section 4 implies that random function  $X$  given by (6.3) is integrable. Thus, it is enough to prove that the equation

$$\int_{\mathbb{R}^d} X(x, t) \varphi(x) dx = \int_{\mathbb{R}^d} A^* \varphi(x) dx \int_0^t X(x, s) ds \quad (6.4)$$

has only the zero solution provided that  $A = A^*$ .

For  $\varphi \in \mathcal{S}(\mathbb{R}^d)$  and  $0 < s < t$  set  $\psi_{t,s}(x) = S(t-s)\varphi(x)$ . Then  $\psi_{t,s} \in \mathcal{S}(\mathbb{R}^d)$ ,  $A\psi_{t,s} + \frac{\partial}{\partial s}\psi_{t,s} = 0$ ,  $\psi_{t,s} \rightarrow \varphi$  uniformly on any bounded subset of  $\mathbb{R}^d$  as  $t \downarrow s$  (see (4.13) [5]), and we get

$$\begin{aligned} \int_{\mathbb{R}^d} X(x,t)\psi_{t,s}(x) dx &\stackrel{(6.4)}{=} \int_{\mathbb{R}^d} A\psi_{t,s}(x) dx \int_0^s X(x,u) du \\ &\stackrel{(5.6)}{=} \int_{\mathbb{R}^d} dx \int_0^s A\psi_{t,u}(x)X(x,u) du + \int_{\mathbb{R}^d} dx \int_0^s A \frac{\partial}{\partial u} \psi_{t,u}(x) du \int_0^u X(x,v) dv \\ &\stackrel{(4.5)}{=} \int_{\mathbb{R}^d} dx \int_0^s A\psi_{t,u}(x)X(x,u) du + \int_0^s du \int_{\mathbb{R}^d} A \frac{\partial}{\partial u} \psi_{t,u}(x) dx \int_0^u X(x,v) dv \\ &\stackrel{(6.4)}{=} \int_{\mathbb{R}^d} dx \int_0^s A\psi_{t,u}(x)X(x,u) du + \int_0^s du \int_{\mathbb{R}^d} \frac{\partial}{\partial u} \psi_{t,u}(x)X(x,u) dx \\ &\stackrel{(4.5)}{=} \int_{\mathbb{R}^d} dx \int_0^s \left( A\psi_{t,u}(x) + \frac{\partial}{\partial u} \psi_{t,u}(x) \right) X(x,u) du = 0. \end{aligned}$$

Passing to the limit as  $t \downarrow s$  and applying Lemma 3.9, we arrive at

$$\int_{\mathbb{R}^d} X(x,s)\varphi(x) dx = 0. \quad \square$$

**Example 6.1.** Let stochastic measure  $\mu$  be generated by a continuous square integrable martingale  $Y$ ,  $\mu(A) = \int_0^T \mathbf{1}_A(t) dY(t)$ ,  $\lambda$  be the Lebesgue measure on  $\mathbb{R}^d$ . Then  $M_t(A) = Y(t)\lambda(A)$ ,  $0 \leq t \leq T$ ,  $A \subset \mathbb{R}^d$ , is a worthy martingale measure with the dominating measure

$$K(A \times B \times (0, t]) = |\langle Y \rangle_t| \lambda(A) \lambda(B)$$

(we use the terminology of [19]). In this case, (6.3) leads to

$$\begin{aligned} X(x,t) &= \int_{\mathbb{R}^d} p(x,y,t)\xi(y) dy + \int_{[0,t]} d\mu(s) \int_{\mathbb{R}^d} p(x,y,t-s)f(y,s) dy \\ &= \int_{\mathbb{R}^d} p(x,y,t)\xi(y) dy + \int_{[0,t] \times \mathbb{R}^d} p(x,y,t-s)f(y,s) M(dy ds). \end{aligned} \tag{6.5}$$

The results of [19, Chapter 2] imply that the integral with respect to  $M(dy ds)$  is well defined and is the limit of integrals of elementary functions. For elementary function, equality of two stochastic integrals in (6.5) is obvious. Further, we can use the dominated convergence theorem for integral with respect to  $d\mu(s)$ .

Similar solution of parabolic SPDE with respect to general martingale measure we have in Example 9 and Remark 20 [3].

**Example 6.2.** Assume that  $\mu$  is generated by real-valued Wiener process  $w$ ,  $\mathcal{J}$  denotes the set of Schwartz rapidly decreasing test functions in  $\mathbb{R}^d$ . Then equation

$$\langle \mathcal{W}(t), \psi \rangle = w(t) \int_{\mathbb{R}^d} \psi(x) dx, \quad \psi \in \mathcal{J},$$

defines the spatially homogeneous Wiener process with values in  $\mathcal{J}'$  (we used the terminology of [10]). For this case, our equality (6.3) is a partial case of mild solution (2.6) [10].

**Remark.** By similar way, we can consider a more general equation

$$dX(x,t) = AX(x,t) dt + \sum_{1 \leq i \leq j} f_i(x,t) d\mu_i(t), \quad X(0) = \xi, \tag{6.6}$$

which includes the case

$$dX(x,t) = AX(x,t) dt + f_1(x,t) dt + f_2(x,t) d\mu(t), \quad X(0) = \xi.$$

The solution of (6.6) is

$$X(x, t) = S(t)\xi(x) + \sum_{1 \leq i \leq j} \int_{[0, t]} [S(t-s)f_i(x, s)] d\mu_i(s).$$

Under assumptions of Theorem 6.1, the solution of (6.6) is unique.

#### REFERENCES

1. S. Albeverio, J.-L. Wu, and T.-S. Zhang, *Parabolic SPDEs driven by Poisson white noise*, Stochastic Process. Appl. **74** (1998), 21–36.
2. G. P. Curbera and O. Delgado, *Optimal domains for  $L_0$ -valued operators via stochastic measures*, Positivity **11** (2007), 399–416.
3. R. C. Dalang, *Extending martingale measure stochastic integral with applications to spatially homogeneous s.p.d.e's*, Electron. J. Probab. **4** (1999), 1–29.
4. G. Da Prato and J. Zabczyk, *Stochastic equations in infinite dimensions*, Encyclopedia Math. Appl., vol. 44, Cambridge Univ. Press, Cambridge, 1992.
5. A. M. Ilyin, A. S. Kalashnikov, and O. A. Oleynik, *Linear second-order partial differential equations of the parabolic type*, J. Math. Sci. (N. Y.) **108** (2002), 435–542.
6. P. Kotelenetz, *Stochastic Ordinary and Stochastic Partial Differential Equations: Transition from Microscopic to Macroscopic Equations*, Stochastic Modelling Appl. Probab., vol. 58, Springer-Verlag, Berlin–Heidelberg–New York, 2007.
7. S. Kwapien and W. A. Woyczyński, *Random Series and Stochastic Integrals: Single and Multiple*, Birkhäuser, Boston, 1992.
8. J. Memin, Yu. Mishura, and E. Valkeila, *Inequalities for the moments of Wiener integrals with respect to a fractional Brownian motion*, Statistics and Probability Letters **27** (2001), 197–206.
9. S. M. Nikolsky, *A Course of Mathematical Analysis*, vol. 2, “Mir”, Moscow, 1977.
10. S. Peszat and J. Zabczyk, *Stochastic evolution equations with a spatially homogeneous Wiener process*, Stochastic Process. Appl. **72** (1997), 187–204.
11. S. Peszat and J. Zabczyk, *Stochastic partial differential equations with Lévy noise: an evolution equation approach*, Encyclopedia Math. Appl., vol. 113, Cambridge Univ. Press, Cambridge, 2007.
12. V. Radchenko, *Integrals with Respect to General Stochastic Measures*, Institute of Mathematics, Kyiv, 1999. (Russian)
13. V. Radchenko, *Heat equation and wave equation with general stochastic measures*, Ukraïn. Mat. Zh. **60** (2008), 1675–1685; English transl. in Ukrainian Math. J. **60** (2008), 1968–1981.
14. V. Radchenko, *Mild solution of the heat equation with a general stochastic measure*, Studia Math. **194** (2009), 231–251.
15. S. Rolewicz, *Metric Linear Spaces*, Monografie Matematyczne, vol. 56, PWN—Polish Scientific Publishers, Warsaw, 1972.
16. C. Ryll-Nardzewski and W. A. Woyczyński, *Bounded multiplier convergence in measure of random vector series*, Proc. Amer. Math. Soc. **53** (1975), 96–98.
17. M. Talagrand, *Les mesures vectorielles à valeurs dans  $L_0$  sont bornées*, Ann. Sci. École Norm. Sup. (4) **14** (1981), 445–452.
18. Ph. Turpin, *Convexités dans les espaces vectoriels topologiques généraux*, Dissertationes Math. **131** (1976).
19. J. B. Walsh, *An introduction to stochastic partial differential equations*, Lect. Not. Math. **1180** (1984), 236–434.

DEPARTMENT OF MATHEMATICAL ANALYSIS, KYIV NATIONAL TARAS SHEVCHENKO UNIVERSITY, KYIV 01601, UKRAINE

*E-mail address:* vradchenko@univ.kiev.ua

Received 22/12/2011